

The deepest radio image using PLoT

By: Jurjen de Jong



Universiteit
Leiden
The Netherlands





SURF





OSCARS
Open Science Clusters' Action
for Research & Society

European LOFAR: 6"






European LOFAR: 0.3"



LoTSS Deep Fields → ELAIS-N1

- ★ 505 hrs of HBA data
- ★ 6'' angular resolution
- ★ 11 $\mu\text{Jy}/\text{beam}$ sensitivity
- ★ Confusion noise!

(Shimwell et al. 2025)



Deeper with LOFAR longest baselines

- ★ International stations -> more collecting area
- ★ Confusion noise negligible at sub-arcsecond resolution

Computational challenges:

- 25 observations -> 400 TB (100 TB compressed)
- 8 billion pixels

Table 1: Metadata from the four ELAIS-N1 observations used in this paper.

Observ. ID	Project	Calibrator ID	Observ. date	Stations (int)	Frequencies
L686892	LT10_012	686888	06-12-2018	70 (12)	120-165 MHz
L686906	LT10_012	L686902	08-12-2018	75 (13)	120-165 MHz
L686920	LT10_012	L686916	03-12-2018	74 (12)	120-165 MHz
L686934	LT10_012	L686940	02-12-2018	74 (12)	120-165 MHz
L686948	LT10_012	L686944	30-11-2018	73 (11)	120-167 MHz
L686962	LT10_012	L686958	26-11-2018	73 (11)	120-165 MHz
L686976	LT10_012	L686972	25-11-2018	73 (11)	128-167 MHz
L687004	LT10_012	L687000	22-11-2018	70 (9)	128-167 MHz
L686990	LT10_012	L686986	24-11-2018	73 (11)	126-167 MHz
L760853	LT10_012	L769403	02-12-2019	67 (12)	120-166 MHz
L769393	LT10_012	L769389	24-05-2020	75 (13)	120-165 MHz
L769407	LT10_012	L769403	08-05-2020	71 (12)	120-165 MHz
L769421	LT10_012	L769417	11-05-2020	73 (13)	120-167 MHz
L769435	LT10_012	L769431	14-03-2020	72 (11)	120-166 MHz
L798022	LT14_003	L798030	12-11-2020	72 (11)	117-166 MHz
L798074	LT14_003	L798082	14-11-2020	74 (11)	115-164 MHz
L798092	LT14_003	L798100	20-11-2020	68 (11)	115-164 MHz
L798128	LT14_003	L798136	26-11-2020	69 (9)	115-166 MHz
L811718	LT14_003	L811726	18-03-2021	72 (11)	117-165 MHz
L811736	LT14_003	L811744	19-03-2021	75 (13)	115-164 MHz
L816272	LT14_003	L816280	13-05-2021	75 (14)	115-164 MHz
L816308	LT14_003	L816316	22-05-2021	75 (13)	116-162 MHz
L816344	LT14_003	L816352	26-05-2021	73 (14)	115-166 MHz
L818954	LT14_003	L818962	21-06-2021	72 (10)	115-156 MHz
L833466	LT14_003	L833474	09-10-2021	73 (13)	119-165 MHz

ELAIS-N1 with 32 hours HBA


★ 4 observations

- 26-11-2018
- 24-05-2020
- 14-11-2020
- 13-5-2021

★ 2 times deeper than 1 observation


★ ~13,000 sources

(See de Jong et al. 2024)






Remaining challenges (de Jong et al. 2024)

- ★ Calibration artefacts (high S/N calibrators)
- ★ Need for more automation
- ★ Imaging cost (85% total compute)




Facet-calibration



Automatic selection

Need enough signal-to-noise
at long baselines




Automatic selection

Need enough signal-to-noise
at long baselines



Metric:

Phase solution difference
between RR-LL polarisation




Validation with AI

- ★ Neural network
- ★ DinoV2 model
- ★ Pre-trained
- ★ 3,000 images




Computational cost (1 observation)




Sidereal visibility averaging

Aperture synthesis





Asaki et al. (2023)


LOFAR UV-coverage



Sidereal visibility averaging (de Jong+2025a)




Computational cost



Comparison with deep ELAIS-N1 6"

(Shimwell et al. 2025)


- ★ **1.8** times deeper
(6 μ Jy/beam !)
- ★ **20 times** better resolution



Comparison with deep ELAIS-N1 6"


(Shimwell et al. 2025)

- ★ 1.8 times deeper
(6 μ Jy/beam !)
- ★ 20 times better resolution




Back to ELAIS-N1

- ★ 32hrs vs. 200hrs
- ★ 2.5 times deeper




Best et al. (2023)




Back to ELAIS-N1


- ★ 32hrs vs. 200hrs
- ★ 2.5 times deeper



Best et al. (2023)



The deepest radio image!



Conclusions

- ★ (Almost) fully automated pipeline
- ★ Computational costs reduced ($\sim 10\times$ in 3 years)
- ★ Deepest ever radio image upcoming!
(Reach out if you want to get early access to the data)

Data processing strategy





Acknowledgement

This presentation is part of the project CORTEX (NWA.1160.18.316) of the research programme NWA-ORC which is (partly) financed by the Dutch Research Council (NWO). This work made use of the Dutch national e-infrastructure with the support of the SURF Cooperative using grant no. EINF-6218. This work is co-funded by the EGI-ACE project (Horizon 2020) under Grant number 101017567.

This work was sponsored by NWO Domain Science for the use of the national computer facilities.



The authors acknowledge the OSCARS project, which has received funding from the European Commission's Horizon Europe Research and Innovation programme under grant agreement No. 101129751

Phasediff-score






Solution interval

$$\sigma_c = \pi \sqrt{1 - \exp\left(-\frac{\zeta}{\sqrt{\delta_t}}\right)} \quad [\text{rad}],$$



Baseline tracks



Nutation, precession, aberration

